

Neutral red retention time assay in determination of toxicity of nanoparticles

Hu, Wentao; Culloty, Sarah; Darmody, Grainne; Lynch, Sharon; Davenport, John; Ramirez-Garcia, Sonia; Dawson, Kenneth; Lynch, Iseult; Doyle, Hugh; Sheehan, David

DOI:

[10.1016/j.marenvres.2015.05.007](https://doi.org/10.1016/j.marenvres.2015.05.007)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Hu, W, Culloty, S, Darmody, G, Lynch, S, Davenport, J, Ramirez-Garcia, S, Dawson, K, Lynch, I, Doyle, H & Sheehan, D 2015, 'Neutral red retention time assay in determination of toxicity of nanoparticles', *Marine Environmental Research*. <https://doi.org/10.1016/j.marenvres.2015.05.007>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published as Hu, W., Culloty, S., Darmody, G., Lynch, S., Davenport, J., Ramirez-Garcia, S., Dawson, K., Lynch, I., Doyle, H., Sheehan, D., Neutral red retention time assay in determination of toxicity of nanoparticles, *Marine Environmental Research* (2015), doi: 10.1016/j.marenvres.2015.05.007. Birmingham

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Neutral red retention time assay in determination of toxicity of nanoparticles

Wentao Hu, Sarah Culloty, Grainne Darmody, Sharon Lynch, John Davenport, Sonia Ramirez-Garcia, Kenneth Dawson, Iseult Lynch, Hugh Doyle, David Sheehan



PII: S0141-1136(15)00074-4

DOI: [10.1016/j.marenvres.2015.05.007](https://doi.org/10.1016/j.marenvres.2015.05.007)

Reference: MERE 4004

To appear in: *Marine Environmental Research*

Received Date: 22 January 2015

Revised Date: 11 May 2015

Accepted Date: 15 May 2015

Please cite this article as: Hu, W., Culloty, S., Darmody, G., Lynch, S., Davenport, J., Ramirez-Garcia, S., Dawson, K., Lynch, I., Doyle, H., Sheehan, D., Neutral red retention time assay in determination of toxicity of nanoparticles, *Marine Environmental Research* (2015), doi: 10.1016/j.marenvres.2015.05.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Neutral red retention time assay in determination of toxicity of nanoparticles

Wentao Hu^a, Sarah Culloty^b, Grainne Darmody^b, Sharon Lynch^b, John Davenport^b,
Sonia Ramirez-Garcia^c, Kenneth Dawson^c, Iseult Lynch^d, Hugh Doyle^e, David
Sheehan^{a*}

^aEnvironmental Research Institute and School of Biochemistry and Cell Biology,
University College Cork, Ireland.

^bAquaculture and Fisheries Development Centre, School of Biological, Earth and
Environmental Sciences, University College Cork, Ireland.

^cCentre for BioNano Interactions and Department of Physical Chemistry, University
College Dublin, Ireland.

^dSchool of Geography, Earth and Environmental Sciences, University of Birmingham,
Edgbaston, Birmingham B 15 2TT, UK.

^eTyndall National Laboratory, University College Cork, Ireland

*Corresponding author's address: School of Biochemistry and Cell Biology, Western
Gateway Building, University College Cork, Cork, Ireland.

Tel: 353 21 4205424

e-mail address: d.sheehan@ucc.ie

Keywords: *Mytilus*, metal oxide, lysosome, membrane stability, neutral red, NRRT

23 **Abstract**

24 The neutral red retention time (NRRT) assay is useful for detecting decreased
25 lysosomal membrane stability in haemocytes sampled from bivalves, a phenomenon
26 often associated with exposure to environmental pollutants including nanomaterials.
27 Bivalves are popular sentinel species in ecotoxicology and use of NRRT in study of
28 species in the genus *Mytilus* is widespread in environmental monitoring. The NRRT
29 assay has been used as an *in vivo* test for toxicity of carbon nanoparticles (Moore MN,
30 Readman JAJ, Readman JW, Lowe DM, Frickers PE, Beesley A. 2009. Lysosomal
31 cytotoxicity of carbon nanoparticles in cells of the molluscan immune system: An *in*
32 *vivo* study. *Nanotoxicology*. 3 (1), 40-45). We here report application of this assay
33 adapted to a microtitre plate format to a panel of metal and metal oxide nanoparticles
34 (2ppm). This showed that copper, chromium and cobalt nanoparticles are toxic by this
35 criterion while gold and titanium nanoparticles are not. As the former three
36 nanoparticles are often reported to be cytotoxic while the latter two are thought to be
37 non-cytotoxic, these data support use of NRRT as a general *in vitro* assay in
38 nanotoxicology.

39

1. Introduction

The unusual properties of nanomaterials provide them with several possible routes to toxicity in biological systems. Their small size sometimes enables them to cross important biobarriers e.g. skin, blood-brain, intestine, maternal-foetus (Tedesco and Sheehan, 2010; Elsaesser and Howard, 2012; Jiang et al., 2014). Their very large surface area to volume ratio enables a greater proportion of atoms to be displayed on the particle surface compared to corresponding macromaterials (Nel et al., 2009; Nel et al., 2013). Moreover, specific functional groups on nanoparticle surfaces may facilitate biospecific interactions allowing a range of possible biological effects (Hoet et al., 2004; Moore, 2006; Klaper et al., 2014). Nanomaterials can also translocate within the human body into other systems such as circulatory and lymphatic vessels (Gwinn and Vallyathan, 2006; Buzea et al., 2007; Elsaesser and Howard, 2012). Thus, nanoparticles have significant potential to cause adverse health effects in humans and other organisms upon prolonged exposure.

Because of increasing commercial production and use of nanomaterials, issues of their accumulation and fate in the environment and their possible effects on ecosystems arise (Moore, 2006; Tedesco and Sheehan, 2010; Ivask et al., 2014). The majority of human habitation worldwide is within 100km of coastlines and the aquatic environment collects domestic, agricultural, shipping and industrial runoffs from these coastal zones. This makes aquatic ecosystems particularly at risk to potential toxicity of nanomaterials of anthropogenic origin. Invertebrates are key elements of the aquatic food chain and mussels are amongst the most abundant of these (Baun et

61 al., 2008). As filter-feeders, mussels are exquisitely selective in the particle size-
62 range which they ingest (Defossez and Hawkins, 1997; Ward and Kach, 2009) and
63 can bioconcentrate metals and organic pollutants within their tissues. This has led to
64 their widespread study in ecotoxicology (Moore, 1985; Widdows and Donkin, 1992)
65 and filter-feeders have been suggested as especially attractive targets for probing the
66 environmental fate of nanomaterials (Moore, 2006; Ward and Kach, 2009; Canesi et
67 al., 2012).

68 Lysosomes are important subcellular organelles that contain many hydrolytic
69 enzymes, carry out protein degradation and detoxify some foreign compounds. At the
70 cellular level, lysosomal digestion pathways include phagocytosis, endocytosis and
71 autophagy. The lysosomal membrane protects the cytosol, and therefore the rest of the
72 cell, from leakage of degradative enzymes. However, malfunctioning of lysosomes
73 and their accumulation of toxic pollutants have been linked to lysosomal storage
74 diseases and result in lysosomal injury and oxidative damage, in some cases leading
75 to cell death (Moore et al., 2007). The neutral red retention time (NRRT) assay takes
76 advantage of this phenomenon by measuring decreased time of retention of a dye,
77 neutral red (ACS no. 553-24-2), within phagocytic haemocytes of a range of aquatic
78 organisms including mussels, crustaceans and fish (Regoli, 1992; Tedesco et al, 2008;
79 Lowe et al 1995; Svendsen et al, 2004). In the popular sentinel species, *Mytilus edulis*,
80 hemocytes are essential immune system components (Rickwood and Galloway,
81 2004). NRTT has been reported as a useful indicator of the organism's overall health

status because animals exposed to pollutants often have compromised lysosomal stability (Moore et al., 2009; Borenfreund and Puerner 1985; Piola et al., 2013). A spectrophotometric version of the assay was developed by Babich and Borenfreund (1990) and a microscopic slide observation method was developed by Moore et al., (2009). This assay takes advantage of the tendency of haemocytes to take up nanoparticles most probably by either phagocytosis or macro-endocytosis and involves exposing haemocytes to nanoparticles on a microscope slide (Moore et al., 2009). In this short report, we have adapted this methodology to a microtitre plate format enabling high-throughput screening of large numbers of replicates, doses and nanoparticles simultaneously (Fig. 1). As proof of principle, we have assessed a panel of metal and metal oxide nanoparticles with this assay.

2. Materials and Methods

2.1. *Mytilus edulis* sampling

M. edulis individuals (4-6cm shell-length) were collected from an intertidal site in Cork Harbour, Ireland (location: 51.49°N, 8 18°W; Lyons et al., 2003). All Animals were acclimated in tanks for a week with a 12 h light/dark cycle at a temperature of 15°C and 34–36‰ salinity, fed and with regular changing of water.

2.2. *Nanoparticle suspension preparation*

Metal or metal oxide nanoparticles (copper oxide, titanium dioxide, gold, chromium oxide and cobalt oxide) of nominal sizes <50nm were purchased from Sigma-Aldrich (Dorset, UK). Nanopowders (10mg) were suspended in 10 ml of 20 mM citric acid adjusted to pH 7, and sonicated for 1h using a tip sonicator. A stepped microtip was used and the total power transferred to the suspension was 2.4W (determined by the calorimetric method). Ultrasound was applied as 15s pulses with 15s breaks between them (Taurozzi et al., 2010). The suspensions were left at 60°C overnight and were then filtered using a 220nm pore size cellulose acetate filter (Millipore, Watford UK).

2.3. *Exposure of haemolymph to nanoparticles*

Haemolymph samples were freshly extracted for NRRT assay as described by Moore et al. (2009). In the present work, haemolymph from each of five animals was extracted from adductor muscle using a 20 gauge hypodermic needle fitted on a 1 ml syringe containing 100µl tris buffered saline buffer, which was pooled to provide a total volume of 2 ml haemolymph solution. Three biologically independent replicates were used (i.e. haemolymph was taken from 3x5 individual animals). Samples were constantly vortexed to resuspend the haemolymph and prevent aggregation. Haemolymph was then evenly aliquoted (500 µL) followed by exposure to nanoparticles at a final concentration of 2 ppm for 1 h at ambient temperature (20°C). Tubes were gently shaken every 5 min to optimise exposure. The above procedure was applied to a panel of metal or metal oxide nanoparticles and a control sample was treated identically but without the presence of nanoparticle.

2.4.Neutral red retention time (NRRT) assay

Following nanoparticle exposure, 100 µl haemolymph from all six treatment groups was loaded into individual wells of a 96-well microtitre plate (Sarstedt, Wexford Ireland). This was performed with three independent biological replicates. Fifty µl stock neutral red dye solution (200 µM) was then added. Four plates were used in parallel for time-points 15, 30, 60 and 90 min. All plates were placed in the dark allowing 15, 30, 60 or 90 min, respectively, for dye uptake. Dye and medium were quickly removed from the plates after incubation and washed with 150 µL

fixative solution (1% formaldehyde, 1% calcium chloride) for 2 min. Plates were then rapidly drained, followed by addition of 200µl extraction buffer (1% acetic acid and 50% ethanol) and left in the dark for 20 min at room temperature. Absorbance of extracted dye was measured using a microplate reader (Elx808iu Ultra Microplate Reader, Bio-Tek Instrument Inc., Potton UK) at a wavelength of 570 nm.

3. Results and Discussion

3.1. Neutral red retention time assay of metal oxide nanoparticles

Haemolymph from *M. edulis* was exposed to a panel of metal or metal oxide nanoparticles at a final concentration of 2ppm (Fig. 1). Lysosomal membrane stability was tested by measuring NRRT at four different time points; 15, 30, 60 and 90 min. Results were analysed and statistically compared to the control group using a one-way anova test with confidence limit of 95% (Figure 2). Lysosomal membrane stability showed a significant decrease ($p < 0.05$) upon exposure to copper, cobalt and chromium nanoparticles at all time-points tested, indicating toxic effects on lysosomes of these nanomaterials. However, no significant effects were observed on exposure of titanium or gold nanoparticles, suggesting they are less toxic by the criterion of this *in vitro* assay.

3.2. Toxicity of metal or metal oxide nanoparticles

The particles selected for this study have previously been reported to display a range of toxicity in biological systems. Titanium dioxide nanoparticles (which are widely used commercially as a component of sunscreens) are generally regarded as less toxic to aquatic species (Federici et al, 2007). However, it should be noted that, in mice, NO and tumour necrosis factor alpha production were elicited after exposure to titanium dioxide nanoparticles (<10nm). This finding suggested that both damage to

the cell structure and macrophage dysfunction may occur, leading to reduction in both non-specific and specific immune responses in some individual animals (Liu et al 2010). Copper oxide and chromium oxide nanoparticles are notorious for their toxic effects, and have been implicated in toxicity to non-target organisms (Ivask et al, 2014), reduction of immune status (Zha et al 2009), damage to animal tissues (Chen et al, 2006; Griffitt et al, 2007), and induction of reactive oxygen species (Fahmy and Cormier, 2009; Horie et al 2011). Cobalt oxide nanoparticles readily enter cultured human cells where they are found to have a negative effect on cell viability (Papis et al., 2009). They have been reported to induce primary DNA damage in a concentration-dependent manner. Various redox enzyme activities were decreased after treatment with cobalt nanoparticles, suggesting potential toxic risk and inhibition of antioxidant capacity (Jiang et al, 2012).

3.3.Potential for high-throughput assay

The assay format reported here includes minimisation of biological variation in haemocyte populations by pooling haemolymph across five individual animals. Moreover, three independent replicates gave essentially identical results and allowed reproducible discrimination across the nanoparticle panel studied. Use of 96-well microtitre plates makes possible high-throughput analysis of large numbers of samples, replicates and concentrations within the time-scale suggested by Moore et al.

(2009). This could facilitate rapid quantitative analysis of novel engineered nanoparticles. An especially attractive feature of this assay format is that it mimics the kinds of strategies that many nanoparticles most probably employ in nature to gain entry to cells such as phagocytosis or macro-endocytosis. This is an ancient and long-established property of eukaryote cells (Elsaesser and Howard, 2012).

Acknowledgement

This study was performed as part of the NeuroNano project (NMP4-SL-2008-214547) funded by the Seventh Framework of the European Union.

References

- Babich, H., Borenfreund, E. 1990. Applications of the neutral red cytotoxicity assay to invitro toxicology. *ATLA - Alt. Lab. Anim.* 18, 129-144.
- Baun, A., Hartmann, N.B., Grieger, K., Kusk, K.O. 2008. Ecotoxicity of engineered nanoparticles to aquatic invertebrates: a brief review and recommendations for future toxicity testing. *Ecotoxicol.* 17, 387-395.
- Borenfreund, E., Puerner, J.A. 1985. Toxicity determined *in vitro* by morphological alterations and neutral red absorption. *Toxicol. Lett.* 24, 119-124.
- Buzea, C., Pacheco, I.I., Robbie, K. 2007. Nanomaterials and nanoparticles: sources and toxicity. *Biointerphases.* 2, 17-71.
- Canesi, L., Ciacci C., Fabbri, R., Marcomini, A., Pojano, G., Gallo, G. 2012. Bivalve molluscs as a unique target group for nanoparticle toxicity. *Mar. Environ. Res.* 76, 16-21.
- Chen, Z., Meng, H., Xiang, G., Chen, C., Zhao, Y., Jia, G., Wang, T., Yuan, H., Ye, C., Zhao, F., Chai, Z., Zhu, C., Fang, X., Ma, B., Wan, L. 2006. Acute toxicological effects of copper nanoparticles *in vivo*. *Toxicol. Lett.* 163, 109-120.
- Defossez, J.M., Hawkins, A.J.S. 1997. Selective feeding in shellfish: Size-dependent rejection of large particles within pseudofeces from *Mytilus edulis*, *Ruditapes philippinarum* and *Tapes decussatus*. *Mar Biol.* 129, 139-147.

- 210 Elsaesser, A., Howard, C.V. 2012. Toxicology of nanoparticles. Adv. Drug Deliv.
211 Rev. 64, 129-137.
- 212 Fahmy, B., Cormier, S.A. 2009. Copper oxide nanoparticles induce oxidative stress
213 and cytotoxicity in airway epithelial cells. Toxicol. in Vitro. 23, 1365-1371.
- Federici, G., Shaw, B.J., Handy, R.D. 2007. Toxicity of titanium dioxide nanoparticles to rainbow trout (*Oncorhynchus mykiss*): Gill injury, oxidative stress, and other physiological effects. Aquat. Toxicol. 84, 415-430.
- Griffitt, R.J., Weil, R., Hyndman, K.A., Denslow, N.D., Powers, K., Taylor, D., Barber, D.S. 2007. Exposure to copper nanoparticles causes gill injury and acute lethality in zebrafish (*Danio rerio*). Environ. Sci. Technol. 41, 8178-8186.
- Gwinn, M.R., Vallyathan, V. 2006. Nanoparticles: health effects--pros and cons. Environ. Health Perspect. 114, 1818-1825.
- 214 Hoet, P.H., Bruske-Holfeld, I., Salata, O.V. 2004. Nanoparticles - known and
215 unknown health risks. J. Nanobiotechnol. 2, 12-15.
- 216 Horie, M., Nishio, K., Endoh, S., Kato, H., Fujita, K., Miyauchi, A., Nakamura, A.,
217 Kinugasa, S., Yamamoto, K., Niki, E., Yoshida, Y., Iwahashi, H. 2011.
218 Chromium(III) oxide nanoparticles induced remarkable oxidative stress and apoptosis
219 on culture cells. Environ. Toxicol. 28, 61-75.
- 220 Ivask, A., Juganson, K., Bondarenko, O., Mortimer, M., Aruoja, V., Kasemets, K.,
221 Blinova, I., Henilaan, M., Slaveykova, V., Kahru, A. 2014. Mechanisms of toxic

- 222 action of Ag, ZnO and CuO nanoparticles to selected ecotoxicological test organisms
223 and mammalian cells *in vitro*: A comparative review. *Nanotoxicol.* 8, 57-71.
- 224 Jiang, C.J., Jia, J.B., Zhai, S.M. 2014. Mechanistic understanding of toxicity from
225 nanocatalysts. *Int. J. Mol. Sci.* 15, 13967-13992.
- 226 Jiang, H., Liu, F., Yang, H., Li, Y. 2012. Effects of cobalt nanoparticles on human T
227 cells *in vitro*. *Biol. Trace Elem. Res.* 146, 23-29.
- 228 Klaper, R., Arndt, D., Bozich, J., Dominguez, G. (2014) Molecular interactions of
229 nanomaterials and organisms: Defining biomarkers for toxicity and high-throughput
230 screening using traditional and next-generation sequencing approaches. *Analyst* 139,
231 882-895.
- 232 Liu, R., Zhang, X., Pu, Y., Yin, L., Li, Y., Zhang, X., Liang, G., Li, X., Zhang, J.
233 2010. Small-sized titanium dioxide nanoparticles mediate immune toxicity in rat
234 pulmonary alveolar macrophages *in vivo*. *J. Nanosci. Nanotechnol.* 10, 5161-5169.
- 235 Lowe, D.M., Fossato, V.U., Depledge, M.H. 1995. Contaminant induced lysosomal
236 membrane damage in blood cells of mussels *M. galloprovincialis* from the Venice
237 Lagoon: An *in vitro* study. *Mar. Ecol. Prog. Ser.* 129, 189-196.
- 238 Lyons, C., Dowling, V., Tedengren, M., Hart, M.G.J., O'Brien, N.M., van Pelt,
239 F.N.A.M., O'Halloran, J., Sheehan, D. 2003. Immunoblotting determination of levels
240 of heat shock protein and glutathione S-transferase in Blue mussel, *Mytilus edulis*,

- 241 sampled from Cork Harbour, Ireland, the North and Baltic Seas. Mar. Environ. Res.
242 56, 585-597.
- 243 Moore, M.N. 1985. Cellular responses to pollutants. Mar. Pollut. Bull.16, 134-139.
- 244 Moore, M.N. 2006. Do nanoparticles present ecotoxicological risks for the health of
245 the aquatic environment? Environ. Internat. 32, 967-976.
- 246 Moore, M.N., Viarengo, A., Donkin, P., Hawkins, A.J.S. 2007. Autophagic and
247 lysosomal reactions to stress in the hepato- pancreas of blue mussels. Aquat. Toxicol.
248 84, 80-91.
- 249 Moore, M.N., Readman, J.A.J., Readman, J.W., Lowe, D.M., Frickers, P.E., Beesley,
250 A. 2009. Lysosomal cytotoxicity of carbon nanoparticles in cells of the molluscan
251 immune system: An *in vivo* study. Nanotoxicol. 3, 40-45.
- 252 Nel, A.E., Madler, L., Velego, D., Xia, T., Hoek, E.M.V., Somosundaran, P.,
253 Klaessig, F., Castranova, V., Thomson, M. 2009. Understanding biophysicochemical
254 interactions at the nano-bio interface. Nat. Mat. 8, 543-557.
- 255 Nel A, Xia T, Meng H, Wang X, Lin SJ, Ji ZX, Zhang HY. 2013. Nanomaterial
256 testing in the 21st Century: Use of a predictive toxicological approach and high-
257 throughput screening. Account. Chem. Res. 46, 607-621.

- 258 Papis, E., Rossi, F., Raspanti, M., Dalle-Donne, I., Colombo, G., Milzani, A.,
259 Bernardinin, G., Gornati, R. 2009. Engineered cobalt oxide nanoparticles readily enter
260 cells. *Toxicol. Lett.* 189, 253-259.
- 261 Piola, L., Fuchs, J., Oneto, M.L., Basack, S., Kesten, E., Casabe, N. (2013)
262 Comparative toxicity of two glyphosate-based formulations to *Eisenia Andrei* under
263 laboratory conditions. *Chemosphere* 91, 545-551.
- 264 Regoli, F. 1992. Lysosomal responses as a sensitive stress index in biomonitoring
265 heavy-metal pollution. *Mar. Ecol. Prog. Ser.* 84, 63-69.
- 266 Rickwood, C.J., Galloway, T.S. 2004. Acetylcholinesterase inhibition as a biomarker
267 of adverse effect: a study of *Mytilus edulis* exposed to the priority pollutant
268 chlorfenvinphos. *Aquat. Toxicol.* 67, 45-56.
- 269 Svendsen, C., Spurgeon, D.J., Hankard, P.K., Weeks, J.M. 2004. A review of
270 lysosomal membrane stability measured by neutral red retention: Is it a workable
271 earthworm biomarker. *Ecotox. Environ. Safe.* 57, 20-29.
- 272 Taurozzi, J.S., Hackley, V.A., Wiesner, M. 2010. Preparation of nanoparticle
273 dispersions from powdered material using ultrasonic disruption. CEINT website. 1-10.
274 http://www.nist.gov/customcf/get_pdf.cfm?pub_id=905633.
- 275 Tedesco, S., Doyle, H., Redmond, G., Sheehan, D. 2008. Gold nanoparticles and
276 oxidative stress in *Mytilus edulis*. *Mar. Environ. Res.* 66, 131-133.

- 277 Tedesco, S., Sheehan, D. 2010. Nanomaterials as Emerging Environmental Threats.
278 Curr. Chem. Biol. 4, 151-160.
- 279 Ward, J.E., Kach, D.J. 2009. Marine aggregates facilitate ingestion of nanoparticles
280 by suspension-feeding bivalves. Mar. Environ. Res. 68, 137-142.
- 281 Widdows, J., Donkin, P. 1992. Mussels and environmental contaminants:
282 bioaccumulation and physiological aspects. In, The Mussel *Mytilus*. Elsevier Press,
283 Amsterdam.
- 284 Zha, L., Zeng, J., Sun, S., Deng, H., Luo, H., Li, W. 2009. Chromium(III)
285 nanoparticles affect hormone and immune responses in heat-stressed rats. Biol. Trace
286 Elem. Res. 129, 157-169.
- 287
- 288

289 **Figure legends**

290 **Figure 1** Schematic overview of NRTT assay.

291 **Figure 2** Neutral red retention time (NRRT) assay in response to a panel of
292 nanoparticles. Neutral red dye extracted from exposed haemocytes was measured
293 spectrophotometrically at 570nm in a plate reader (*p< 0.05 versus control values).

294

295

296

297




Fig. 1




Figure 1

Figure 2

- Neutral red retention time assay used haemolymph of five pooled mussels.
- Assay was miniaturised for reading in a plate reader, facilitating many samples and replicates.
- Copper, chromium and cobalt nanoparticles were toxic while gold and titanium were not.